

Anti-HSP90 complex

Catalog# SMC-109A/B

Size: 50/200µg

PO Box 30244, Suite 405,
3989 Quadra Street,
Victoria, BC V8X 5E1, Canada

This product is for *in vitro* research use only and is not intended for use in humans or animals

StressMarq

Biosciences Inc.

Orders ● sales@stressmarq.com
Tel: ● +1 250 294 9065
Fax: ● +1 250 294 9025
Email ● info@stressmarq.com
Web ● www.stressmarq.com

Product	Mouse anti-hsp90, 8D3 monoclonal antibody
Clone	8D3
Immunogen	Ah receptor (Aryl hydrocarbon receptor) (8)
Host and Subclass	Mouse IgM
Applications	IP
Specificity	Immunoprecipitates 90kD proteins corresponding to the molecular mass of hsp90. Co-immunoprecipitates hsp90 complexes, including hsp70, Hop, Ah receptors (8), glucocorticoid receptors (9), heme-regulated eukaryotic initiation factor 2alpha (eIF-2alpha) kinase (HRI) (10)
Species cross-reactivity	Human, Mouse, Rat, Rabbit. Not tested on other species.
Format	In PBS in 0.09% sodium azide in 50% glycerol. PEG purified.
Concentration and working dilution	1.0mg/mL; best results for IP associated with use of goat anti-mouse IgM beads.
Storage and stability	-20°C; 1 year+; shipped on cold packs or ambient

Scientific Background

Hsp90 is a highly conserved and essential stress protein that is expressed in all eukaryotic cells. From a functional perspective, hsp90 participates in the folding, assembly, maturation, and stabilization of specific proteins as an integral component of a chaperone complex (1-4). Despite its label of being a heat-shock protein, hsp90 is one of the most highly expressed proteins in unstressed cells (1-2% of cytosolic protein). It carries out a number of housekeeping functions - including controlling the activity, turnover, and trafficking of a variety of proteins. Most of the hsp90-regulated proteins that have been discovered to date are

involved in cell signaling (5-6). The number of proteins now known to interact with Hsp90 is about 100. Target proteins include the kinases v-Src, Wee1, and c-Raf, transcriptional regulators such as p53 and steroid receptors, and the polymerases of the hepatitis B virus and telomerase.5 When bound to ATP, Hsp90 interacts with co-chaperones Cdc37, p23, and an assortment of immunophilin-like proteins, forming a complex that stabilizes and protects target proteins from proteasomal degradation.

In most cases, hsp90-interacting proteins have been shown to co-precipitate with hsp90 when carrying out immunoadsorption studies, and to exist in cytosolic heterocomplexes with it. In a number of cases, variations in hsp90 expression or hsp90 mutation has been shown to degrade signaling function via the protein or to impair a specific function of the protein (such as steroid binding, kinase activity) *in vivo*. Ansamycin antibiotics, such as geldanamycin and radicicol, inhibit hsp90 function (7).

Selected References

1. Arlander SJH, et al. (2003) *J Biol Chem* 278: 52572-52577.
2. Pearl H, et al. (2001) *Adv Protein Chem* 59:157-186.
3. Neckers L, et al. (2002) *Trends Mol Med* 8:S55-S61.
4. Pratt W, Toft D. (2003) *Exp Biol Med* 228:111-133.
5. Pratt W, Toft D. (1997) *Endocr Rev* 18: 306-360.
6. Pratt WB. (1998) *Proc Soc Exptl Biol Med* 217: 420-434.
7. Whitesell L, et al. (1994) *Proc Natl Acad Sci USA* 91: 8324-8328.
8. Perdew, G. H. (1988) *JBC* 263 (27): 13802-13805
9. Dalman, F. C. et al. (1989) *JBC* 264(33): 19815-19821
10. Uma, S. et al. (1997) *JBC* 272(17): 11648-11656

Certificate of Analysis

Goat anti-mouse IgM was used to bind 25 µl of protein G-Sepharose. SMC-109 IgM from 0.5 ml of high speed supernatant medium was loaded onto the IgG resin and incubated with 100 µl of rabbit reticulocyte lysate for 30 min. at 30C. After washing (4X1 ml), bound proteins were resolved on SDS PAGE, including hsp90, hsp70 and Hop.

Material Safety Data Sheet

Anti-Hsp90, 8D3 (Monoclonal Antibody) SMC-109

This product is for *in vitro* research use only and is not intended for use in humans or animals

The below information is believed to be correct but does not purport to be all inclusive and shall be used only as a guide. StressMarq shall not be held liable for any damage resulting from handling or from contact with the above product. See the Technical Specification, Packing Slip, Invoice, and Product Catalogue for additional terms and conditions of sale.

Hazardous Ingredients

The physical, chemical and toxicological properties of these components have not been fully investigated. It is recommended that all laboratory personnel follow standard laboratory safety procedures when handling this product. Safety procedures should include wearing OSHA approved safety glasses, gloves and protective clothing. Direct physical contact with this product should be avoided.

<u>Known Hazardous Components</u>	<u>CAS Number</u>	<u>Percent</u>
Sodium Azide	26628-22-8	0.09

Physical Data

This product consists of mouse immunoglobulin in PBS containing 0.09% azide in 50% glycerol shipped on gel packs. The physical properties of this product have not been investigated thoroughly.

Fire and Explosion Hazard and Reactivity Data

NOT APPLICABLE

Toxicological Properties

May be harmful by inhalation, ingestion, or skin absorption. The toxicological properties of this product have not been investigated thoroughly. Exercise due caution.

Preventative Measures

Wear chemical safety goggles and compatible chemical-resistant gloves. Avoid inhalation, contact with eyes, skin or clothing.

Spill and Leak Procedures

Observe all federal, state and local environmental regulations.

- Wear protective equipment.
- Absorb on sand or vermiculite and place in closed containers for disposal.
- Dispose or mix the material with a combustible solvent and burn in a chemical incinerator equipped with an afterburner and scrubber.

First Aid Measures

- If swallowed, wash out mouth with water, provided person is conscious. Call a physician.
- In case of skin contact, flush with copious amounts of water for at least 15 minutes. Remove contaminated clothing and shoes. If a rash or other irritation develops, call a physician.
- If inhaled, remove to fresh air. If breathing becomes difficult, call a physician.
- In case of eye contact, flush with copious amounts of water for at least 15 minutes while separating the eyelids with fingers. Call a physician.

Authorized: StressMarq Biosciences Inc.
Creation Date: 12/17/07